

Roll No. 

Total No. of Pages : 03

Total No. of Questions : 09

**B.Tech. (2011 Onwards) (Sem.-1)**  
**ENGINEERING MATHEMATICS – I**  
 Subject Code : BTAM-101  
 Paper ID : [A1101]

Time : 3 Hrs.

Max. Marks : 60

**INSTRUCTIONS TO CANDIDATES :**

1. SECTION-A is COMPULSORY consisting of TEN questions carrying TWO marks each.
2. SECTION - B & C. have FOUR questions each.
3. Attempt any FIVE questions from SECTION B & C carrying EIGHT marks each.
4. Select atleast TWO questions from SECTION - B & C.
5. Symbols used have their usual meanings. Statistical tables, if demanded, may be provided.

**SECTION-A**

1. Solve the following :

a) If  $u = f\left(\frac{y-x}{xy}, \frac{z-x}{xz}\right)$ , then show that  $x^2 \frac{\partial u}{\partial x} + y^2 \frac{\partial u}{\partial y} + z^2 \frac{\partial u}{\partial z} = 0$ .

b) Find the stationary points of the function  $f(x, y) = x^3 + y^3 - 63(x + y) + 12xy$ .

c) If  $u = x^2 - y^2$  and  $v = 2xy$  and  $x = r \cos \theta$ ,  $y = r \sin \theta$ , then find the value of  $\frac{\partial(u, v)}{\partial(r, \theta)}$ .

d) For what values of  $a$ ,  $b$ , and  $c$  the vector function

$$\vec{F} = (x + y + az)\vec{i} + (bx + 3y - z)\vec{j} + (3x + cy + z)\vec{k} \text{ is irrotational.}$$

e) Calculate the circulation of the field  $\vec{F} = (x - y)\hat{i} + x\hat{j}$  around the circle  $x^2 + y^2 = 1$ .

f) Find the length of one arc of the cycloid  $x = a(\theta - \sin \theta)$ ,  $y = a(1 - \cos \theta)$ .

g) Evaluate  $\int_0^{2\sqrt{\ln 3}} \int_{y/2}^{\sqrt{\ln 3}} e^{x^2} dx dy$ .

h) State Stoke's theorem.

i) Evaluate  $\int_C xy dx + (x + y) dy$ , along the curve  $C : y = x^2$  from  $(-1, 1)$  to  $(2, 4)$ .

j) Evaluate :  $\int_0^a \int_0^x \int_0^{x+y} e^{x+y+z} dz dy dx$ .

## SECTION-B

2. a) Find the radius of curvature at any point of the curve  $r = a(1 - \cos \theta)$  and prove that  $\rho^2 / r$  is constant.
- b) Trace the curve  $y^2(x^2 + y^2) + a^2(x^2 - y^2) = 0$  by giving all its features in detail.
3. a) Find the volume of the solid formed by revolving the curve  $y^2(2a - x) = x^3$  about its asymptote.
- b) Find the centre of gravity of the arc of the curve  $x = a(t + \sin t)$ ,  $y = a(1 - \cos t)$  in the first quadrant.
4. a) If  $u = f(r)$ , where  $r^2 = x^2 + y^2$ , then show that  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = f''(r) + \frac{1}{r} f'(r)$
- b) Use Lagrange's method of undetermined coefficients to show that the rectangular solid of maximum volume that can be inscribed in a sphere is a cube.
5. a) Find the first three terms of the Taylor's series expansion of  $e^x \log(1 + y)$  in the neighbourhood of  $(0,0)$
- b) Use Euler's theorem to prove that

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = \frac{\sin u \cos 2u}{4 \cos^3 u}, \text{ whenever } u = \sin^{-1} \left( \frac{x+y}{\sqrt{x} + \sqrt{y}} \right)$$

## SECTION-C

6. a) Evaluate  $\int_0^{2\sqrt{2x-x^2}} \int_0^x (x^2 + y^2) dy dx$  by changing into polar coordinates.
- b) Find the volume bounded by the cylinder  $x^2 + y^2 = 4$  and the planes  $y + z = 4$  and  $z = 0$ .
7. a) Prove the identity :

$$\nabla \left[ \frac{\vec{a} \cdot \vec{r}}{r^n} \right] = \frac{\vec{a}}{r^n} - \frac{n(\vec{a} \cdot \vec{r}) \vec{r}}{r^{n+2}}, \text{ where } \vec{a} \text{ is a constant vector.}$$

b) A vector field is given by  $\vec{F} = 18z\hat{i} - 12\hat{j} + 3y\hat{k}$ . Evaluate the surface integral

$$\int_S \vec{F} \cdot \hat{n} ds, \text{ where } S \text{ is the part of the plane } 2x + 3y + 6z = 12 \text{ in the first octant.}$$

8. Verify the Gauss Divergence theorem for a vector field defined by  $\vec{F} = (x^2 - yz)\hat{i} + (y^2 - xz)\hat{j} + (z^2 - xy)\hat{k}$  taken around the rectangular parallelepiped  $0 \leq x \leq a, 0 \leq y \leq b, 0 \leq z \leq c$ .

9. a) Find the directional derivative of  $f(x, y, z) = x y^2 + y z^3$  at  $(2, -1, 1)$  in the direction of normal to the surface  $x \log z - y^2 = -4$  at  $(-1, 2, 1)$ .

b) State Green's theorem in plane and use it to evaluate

$$\int_C (3x^2 - 8y^2) dx + (4y - 6xy) dy,$$

where C is the boundary of the region defined by  $x = 0, y = 0, x + y = 1$